



Error estimates for some composite corrected quadrature rules

Zheng Liu

Institute of Applied Mathematics, School of Science, University of Science and Technology Liaoning, Anshan 114051, Liaoning, China

ARTICLE INFO

Article history:

Received 3 November 2006

Received in revised form 25 June 2008

Accepted 26 August 2008

Keywords:

Corrected trapezoidal rule

Corrected midpoint rule

Corrected Simpson rule

Composite corrected quadrature rule

Error estimates

ABSTRACT

The asymptotic behaviour of the error for a general quadrature rule is established and it is applied to some composite corrected quadrature rules.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

Let $f : [a, b] \rightarrow \mathbf{R}$ be such that f' is absolutely continuous on $[a, b]$. It is well known that the trapezoidal rule

$$T(f) := \frac{b-a}{2}[f(a) + f(b)] \quad (1)$$

and the midpoint rule

$$M(f) := (b-a)f\left(\frac{a+b}{2}\right) \quad (2)$$

are the simplest quadrature formulae used to approximate the integral

$$\int_a^b f(t)dt$$

which can serve as basic elements for constructing more sophisticated formulae by certain types of convex combinations, e.g., the classical Simpson rule is defined as

$$S(f) = \frac{1}{3}T(f) + \frac{2}{3}M(f) = \frac{b-a}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right]. \quad (3)$$

Further, we may consider the corrected or perturbed trapezoidal rule

$$CT(f) := \frac{b-a}{2}[f(a) + f(b)] - \frac{(b-a)^2}{12}[f'(b) - f'(a)] \quad (4)$$

E-mail address: lewzheng@163.net.

and the corrected or perturbed midpoint rule

$$CM(f) := (b-a)f\left(\frac{a+b}{2}\right) + \frac{(b-a)^2}{24}[f'(b) - f'(a)]. \quad (5)$$

It is well known that corrected quadrature formulae (4) and (5) have better estimates of error than corresponding original formulae (1) and (2). (See, e.g., [1–3].)

Unfortunately, we could not obtain such a corrected version of the Simpson rule, since

$$\frac{1}{3}CT(f) + \frac{2}{3}CM(f) = \frac{b-a}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right]$$

does not involve the derivatives at the endpoints.

Nevertheless, we did find in [4–6] the so-called corrected Simpson quadrature rule that improves on Simpson rule (3):

$$\begin{aligned} CS(f) &= \frac{7}{15}CT(f) + \frac{8}{15}CM(f) \\ &= \frac{b-a}{30} \left[7f(a) + 16f\left(\frac{a+b}{2}\right) + 7f(b) \right] - \frac{(b-a)^2}{60}[f'(b) - f'(a)]. \end{aligned} \quad (6)$$

It is not difficult to find that the corrected trapezoidal rule (4) and the corrected midpoint rule (5) are exact for polynomials of degree 3 or less, and the corrected Simpson rule (6) is exact for polynomials of degree 5 or less.

Now let $a = t_0 < t_1 < \dots < t_n = b$ be an equidistant subdivision of the interval $[a, b]$ such that $t_{i+1} - t_i = h = \frac{b-a}{n}$, $i = 0, 1, \dots, n-1$. Then we have the composite corrected trapezoidal rule

$$CT_n(f) = \frac{b-a}{2n} [f(t_0) + 2f(t_1) + \dots + 2f(t_{n-1}) + f(t_n)] - \frac{(b-a)^2}{12n^2} [f'(b) - f'(a)], \quad (7)$$

the composite corrected midpoint rule

$$CM_n(f) = \frac{b-a}{n} \left[f\left(\frac{t_0+t_1}{2}\right) + f\left(\frac{t_1+t_2}{2}\right) + \dots + f\left(\frac{t_{n-1}+t_n}{2}\right) \right] + \frac{(b-a)^2}{24n^2} [f'(b) - f'(a)] \quad (8)$$

and the composite corrected Simpson rule

$$\begin{aligned} CS_n(f) &= \frac{b-a}{30n} \left\{ 7[f(t_0) + 2f(t_1) + \dots + 2f(t_{n-1}) + f(t_n)] \right. \\ &\quad \left. + 16 \left[f\left(\frac{t_0+t_1}{2}\right) + f\left(\frac{t_1+t_2}{2}\right) + \dots + f\left(\frac{t_{n-1}+t_n}{2}\right) \right] \right\} - \frac{(b-a)^2}{60n^2} [f'(b) - f'(a)] \end{aligned} \quad (9)$$

which correspond to the corrected quadrature rules (4)–(6), respectively.

Motivated by [7,8], in this work, we will give a unified treatment for estimating the errors for the above mentioned composite corrected quadrature rules (7)–(9).

2. Main results

We need the following result (see Theorem 10 in [8] or Theorem 3 in [9]):

Lemma 1. Let $f : [a, b] \rightarrow \mathbf{R}$ be such that its $(r-1)$ th derivative $f^{(r-1)}$ is of continuous bounded variation for some positive integer r . Then for any $x \in [0, 1]$, we have

$$\int_a^b f(t)dt = h \sum_{i=0}^{n-1} f(t_i + xh) - \sum_{v=1}^r \frac{f^{(v-1)}(b) - f^{(v-1)}(a)}{v!} B_v(x)h^v + R_n^{(r)}(f; x), \quad (10)$$

where

$$R_n^{(r)}(f; x) = \frac{h^r}{r!} \int_a^b \tilde{B}_r \left(x - n \frac{t-a}{b-a} \right) df^{(r-1)}(t) \quad (11)$$

and $\tilde{B}_r(t) := B_r(t - [t])$ while $B_r(t)$ is the r th Bernoulli polynomial.

Theorem 2. Suppose that for some real constants $\alpha \neq 0$ and $p_j \in [0, 1], j = 0, 1, \dots, m-1$, with $\sum_{j=0}^{m-1} p_j = 1$, the following quadrature rule

$$\int_0^1 f(t)dt = \sum_{j=0}^{m-1} p_j f(x_j) + \alpha[f'(1) - f'(0)] \quad (12)$$

is exact for any polynomial of degree $\leq r-1$ for some positive integer $r \geq 3$. Let $f : [a, b] \rightarrow \mathbf{R}$ be such that its $(r-1)$ th derivative $f^{(r-1)}$ is a continuous function of bounded variation on $[a, b]$. Then we have

$$\int_a^b f(t)dt = h \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} p_j f(t_i + x_j h) + \alpha h^2 [f'(b) - f'(a)] + R_n^{(r)}(f), \quad (13)$$

where

$$R_n^{(r)}(f) = h^r \int_a^b G_r \left(n \frac{t-a}{b-a} \right) df^{(r-1)}(t), \quad (14)$$

and

$$G_r(t) = \frac{1}{r!} \sum_{j=0}^{m-1} p_j (\tilde{B}_r(x_j - t) - B_r(x_j)). \quad (15)$$

Proof. We first note that for $1 \leq v \leq r-1$,

$$\sum_{j=0}^{m-1} p_j B_v(x_j) + \alpha[B'_v(1) - B'_v(0)] = \int_0^1 B_v(t)dt = 0$$

due to (12) being exact for any polynomial of degree $\leq r-1$ and a well known property of the Bernoulli polynomials. Moreover, since $B_0(t) = 1$, $B'_n(t) = nB_{n-1}(t)$ and $B_n(t+1) - B_n(t) = nt^{n-1}$, $n = 1, 2, \dots$, we get

$$\sum_{j=0}^{m-1} p_j B_v(x_j) = \begin{cases} 0, & \text{if } 1 \leq v \leq r-1, v \neq 2, \\ -2\alpha, & \text{if } v = 2. \end{cases} \quad (16)$$

Now setting $x = x_j$ in (10), multiplying both sides of (10) by p_j , summing from $j = 0$ to $m-1$, from (10), (16) and the following readily checked fact:

$$\frac{f^{(r-1)}(b) - f^{(r-1)}(a)}{r!} B_r(x) h^r = \frac{h^r}{r!} \int_a^b B_r(x) df^{(r-1)}(t),$$

we obtain

$$\int_a^b f(t)dt = h \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} p_j f(t_i + x_j h) - \sum_{v=1}^r \left(\sum_{j=0}^{m-1} p_j B_v(x_j) \right) \frac{f^{(v-1)}(b) - f^{(v-1)}(a)}{v!} h^v + \sum_{j=0}^{m-1} p_j R_n^{(r)}(f; x_j),$$

i.e.,

$$\begin{aligned} \int_a^b f(t)dt &= h \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} p_j f(t_i + x_j h) - \sum_{v=1}^{r-1} \left(\sum_{j=0}^{m-1} p_j B_v(x_j) \right) \frac{f^{(v-1)}(b) - f^{(v-1)}(a)}{v!} h^v \\ &\quad - \sum_{j=0}^{m-1} p_j B_r(x_j) \frac{f^{(r-1)}(b) - f^{(r-1)}(a)}{r!} h^r + \sum_{j=0}^{m-1} p_j R_n^{(r)}(f; x_j), \end{aligned}$$

and it follows from (16) that

$$\begin{aligned} \int_a^b f(t)dt &= h \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} p_j f(t_i + x_j h) + \alpha h^2 [f'(b) - f'(a)] \\ &\quad - \sum_{j=0}^{m-1} p_j \frac{h^r}{r!} \int_a^b B_r(x_j) df^{(r-1)}(t) + \sum_{j=0}^{m-1} p_j R_n^{(r)}(f; x_j). \end{aligned}$$

Finally, from (11) we get

$$\int_a^b f(t)dt = h \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} p_j f(t_i + x_j h) + \alpha h^2 [f'(b) - f'(a)] + \frac{h^r}{r!} \int_a^b \sum_{j=0}^{m-1} p_j (\tilde{B}_r \left(x_j - n \frac{t-a}{b-a} \right) - B_r(x_j)) df^{(r-1)}(t),$$

which completes the proof. \square

Remark 3. The case $\alpha = 0$ has already been studied in [8]. For $\alpha \neq 0$, from the proof of Theorem 2, we can conclude that

$$\sum_{v=1}^{r-1} \left(\sum_{j=0}^{m-1} p_j B_v(x_j) \right) \frac{f^{(v-1)}(b) - f^{(v-1)}(a)}{v!} h^v = -\alpha h^2 [f'(b) - f'(a)]$$

only if $r \geq 3$. Otherwise (for $r = 1, 2$) the expression on the left-hand side equals 0.

Theorem 4. Let the assumptions of Theorem 2 hold. Then we have

$$|n^r R_n^{(r)}(f)| \leq C_r (b-a)^r \bigvee_a^b(f^{(r-1)}), \quad (17)$$

where

$$C_r := \frac{1}{r!} \sup_{0 < t < 1} \left| \sum_{j=0}^{m-1} p_j (\tilde{B}_r(x_j - t) - B_r(x_j)) \right|. \quad (18)$$

If further $f^{(r-1)}$ is absolutely continuous on $[a, b]$, we then have

$$\lim_{n \rightarrow \infty} n^r R_n^{(r)}(f) = K_r (b-a)^r \int_a^b f^{(r)}(t) dt, \quad (19)$$

where

$$K_r := -\frac{1}{r!} \sum_{j=0}^{m-1} p_j B_r(x_j). \quad (20)$$

The proof is similar to that for the Theorem 12 in [8] and so is omitted.

Remark 5. If (12) has degree of precision $r-1$ then C_v and K_v exist for all $3 \leq v \leq r$ since in this case (12) is exact for any polynomial of degree $\leq v-1$. We obtain from (16) and (20) that

$$K_v = 0, \quad \text{if } 3 \leq v < r, \quad (21)$$

which implies from (19) that

$$\lim_{n \rightarrow \infty} n^v R_n^{(v)}(f) = 0, \quad \text{if } 3 \leq v < r.$$

3. Examples

Example 1. For the corrected trapezoid rule, we see that $\alpha = -\frac{1}{12}$ and it has degree of precision 3 ($r = 4$). Thus for any function f such that $f^{(v-1)}$ ($v = 3, 4$) is continuous of bounded variation, we get C_v by direct calculations from (18) as

$$C_3 = \frac{\sqrt{3}}{216}, \quad C_4 = \frac{1}{384},$$

and, for any function f such that $f^{(v-1)}$ ($v = 3, 4$) is absolutely continuous, we obtain K_v from (21) and by a direct calculation from (20) as

$$K_3 = 0, \quad K_4 = \frac{1}{720}$$

and these imply that

$$\lim_{n \rightarrow \infty} n^3 R(f; CT_n) = \lim_{n \rightarrow \infty} n^3 R_n^{(3)}(f) = 0$$

and

$$\lim_{n \rightarrow \infty} n^4 R(f; CT_n) = \lim_{n \rightarrow \infty} n^4 R_n^{(4)}(f) = \frac{1}{720} (b-a)^4 [f'''(b) - f'''(a)].$$

Example 2. For the corrected midpoint rule, we see that $\alpha = \frac{1}{24}$ and it has degree of precision 3 ($r = 4$). Thus for any function f such that $f^{(\nu-1)}$ ($\nu = 3, 4$) is continuous of bounded variation, we get C_ν by direct calculations from (18) as

$$C_3 = \frac{\sqrt{3}}{216}, \quad C_4 = \frac{1}{384},$$

and, for any function f such that $f^{(\nu-1)}$ ($\nu = 3, 4$) is absolutely continuous, we obtain K_ν from (21) and by a direct calculation from (20) as

$$K_3 = 0, \quad K_4 = -\frac{7}{5760}$$

and these imply that

$$\lim_{n \rightarrow \infty} n^3 R(f; CM_n) = \lim_{n \rightarrow \infty} n^3 R_n^{(3)}(f) = 0$$

and

$$\lim_{n \rightarrow \infty} n^4 R(f; CM_n) = \lim_{n \rightarrow \infty} n^4 R_n^{(4)}(f) = -\frac{7}{5760} (b-a)^4 [f'''(b) - f'''(a)].$$

Example 3. For the corrected Simpson rule, we see that $\alpha = -\frac{1}{60}$ and it has degree of precision 5 ($r = 6$). Thus for any function f such that $f^{(\nu-1)}$ ($\nu = 3, 4, 5, 6$) is continuous of bounded variation, we get C_ν by direct calculations from (18) as

$$C_3 = \frac{7}{20250} + \frac{19\sqrt{19}}{81000}, \quad C_4 = \frac{1}{5760}, \quad C_5 = \frac{1}{58320}, \quad C_6 = \frac{1}{230400}$$

and, for any function f such that $f^{(\nu-1)}$ ($\nu = 3, 4, 5, 6$) is absolutely continuous, we obtain K_ν from (21) and by a direct calculation from (20) as

$$K_3 = K_4 = K_5 = 0, \quad K_6 = \frac{1}{604800}$$

and these imply that

$$\lim_{n \rightarrow \infty} n^\nu R(f; CS_n) = \lim_{n \rightarrow \infty} n^\nu R_n^{(\nu)}(f) = 0, \quad \text{for } \nu = 3, 4, 5,$$

and

$$\lim_{n \rightarrow \infty} n^6 R(f; CS_n) = \lim_{n \rightarrow \infty} n^6 R_n^{(6)}(f) = \frac{1}{604800} (b-a)^4 [f^{(5)}(b) - f^{(5)}(a)].$$

Remark 6. It should be mentioned that the same estimates of error for the corrected Simpson rule were obtained in [6]. Furthermore, estimates of error for both classical and corrected trapezoidal rules were obtained in [10] and for the midpoint and corrected midpoint rules in [11].

Acknowledgment

The author wishes to express his sincere thanks to the referee for valuable comments and suggestions.

References

- [1] P. Cerone, On perturbed trapezoidal and midpoint rules, Korean J. Comput. Appl. Math. 2 (2002) 423–435.
- [2] X.L. Cheng, J. Sun, A note on the perturbed trapezoid inequality, J. Inequal. Pure Appl. Math. 3 (2002). Article 29 <http://jipam.vu.edu.au>.
- [3] Z. Liu, On sharp perturbed midpoint inequalities, Tamkang J. Math. 36 (2005) 131–136.
- [4] N. Ujević, A.J. Roberts, A corrected quadrature formula and applications, ANZIAM J. 45 (E) (2004) E41–E56.
- [5] N. Ujević, A generalization of the modified Simpson's rule and error bounds, ANZIAM J. 47 (E) (2005) E1–E13.
- [6] J. Pečarić, I. Franjić, Generalization of corrected Simpson's formula, ANZIAM J. 47 (2006) 367–385.
- [7] D. Cruz-Uribe, C.J. Neugebauer, Sharp error bounds for the trapezoidal rule and Simpson's rule, J. Inequal. Pure Appl. Math. 3 (2002). Article 49 <http://jipam.vu.edu.au>.
- [8] Q.B. Wu, S.J. Yang, A note to Ujević's generalization of Ostrowski's inequality, Appl. Math. Lett. 18 (2005) 657–665.
- [9] I. Franjić, I. Perić, J. Pečarić, General Euler–Ostrowski formulae and applications to quadratures, Appl. Math. Comput. 177 (2006) 92–98.
- [10] Lj. Dedić, M. Matić, J. Pečarić, On Euler trapezoid formulae, Appl. Math. Comput. 123 (2001) 37–62.
- [11] Lj. Dedić, M. Matić, J. Pečarić, On Euler midpoint formulae, ANZIAM J. 46 (2005) 417–438.